




Strategic Research Agenda of the Multidisciplinary European Low Dose Initiative (MELODI) – 2026

F. Antonelli, A. Auvinen, C. Badie, G. Baiocco, S. Candéias, U. Galderisi, L. Lundholm, B. Madas, S. Moertl, R. Quintens, Y. Saintigny.

S. Haghdoost (SRA working group observer)

MELODI Working Group SRA

Francesca Antonelli, ENEA, Italy

Anssi Auvinen, University of Tampere, STUK, Finland

Christophe Badie, UKHSA

Giorgio Baiocco (vice chair), University of Pavia, Italy

Serge Candéias, CEA, France

Yannick Saintigny, CEA, France

Umberto Galderisi, University of Campania

Lovisa Lundholm, Stockholm University/Swedish Radiation Safety Authority, Sweden

Balázs Madas (chair), HUN-REN EK, Hungary

Simone Moertl, BfS, Germany

Roel Quintens, SCK CEN, Belgium

Observers from other MELODI WGs

Siamak Haghdoost, University of Caen, France (Education & Training WG)

Contents

1. Executive Summary	4
2. Importance of low dose radiation health risk research.....	7
2.1 Dose and dose rate ranges to be considered	8
3. MELODI	9
4. List of recent projects	11
5. Strategic Research Agenda	17
5.1 Dose and dose-rate dependence of cancer risk	19
5.2 Non-cancer effects.....	23
5.3 Individual variation in risk.....	26
5.4 Effects of spatial and temporal variation in dose delivery	28
6. Education and Training	31
6.1 The role of education and training in low-dose radiation research	31
6.2 Priorities for strategic support of E&T	32
7. Infrastructures	33
8. Abbreviations, Websites.....	35
9. Annex: Publications from MELODI Workshops	35

1. Executive Summary

The Multidisciplinary European Low Dose Initiative (MELODI) is a European Platform dedicated to research on risks from low-dose ionizing radiation. The challenge is to improve the quantification of risks and reduce the uncertainties in the risk estimates, as well as to develop and validate risk models that best characterise health effects at low doses, drawing on both epidemiological and radiobiological understanding. In 2010, MELODI was founded as a registered association with 15 members; now it has 37 member institutions.

A major activity of MELODI is the establishment and periodic revision of a long-term Strategic Research Agenda (SRA) for research on low-dose risk for radiation protection (RP) in Europe. MELODI considers low doses to be those where there remains substantial uncertainty on the magnitude of health risk. The SRA is intended to guide the priorities for national and European research programmes and the preparation of competitive calls at the European level in order to fill research gaps and test the hypotheses on which the current RP system is based. The ultimate goal is to provide improved evidence-based protection of the population. A key priority for radiation protection research is to reduce the uncertainties associated with health risk estimates for exposures at low doses and dose rates that are relevant for the dose limits for occupational exposures, reference levels for the exposure of the population in emergency situations, diagnostic reference levels for medical exposures, damage to normal tissues during radiotherapy, reference levels for radon exposures in buildings and occupational compensation scheme claims, amongst others. The approaches have to be multidisciplinary and innovative to provide the best opportunities for advancing the understanding of low dose and low dose-rate effects. Incorporation of expertise outside of the conventional fields of radiation research is essential to widen the prospects for broadening approaches and adopting novel methods in the assessment of health risk relevant to radiation protection. MELODI also aims to ensure the availability of key infrastructures as an essential basis for research activities, and to maintain competences in radiation protection research and health risk assessment in the long term via an integrated European approach for training and education. For these purposes, MELODI established three working groups (WGs) in February 2014, one of them is the MELODI SRA WG.

The SRA is periodically updated by the MELODI SRA Working Group (WG), systematically taking into account results of recent research and emerging radiation protection research issues. Open consultations via the website and the annual MELODI workshops are regularly conducted, the results of which are taken into account in the revised SRA.

Large parts of radiation protection research in Europe have been organized within a European Joint Programme (EJP), CONCERT (2015-2020). The aim of the EJP was to bring together relevant funding agencies from the EC and its Member States to integrate European research and to administer calls for research proposals in radiation protection on behalf of the EC. This activity is built upon to promote integration of the SRAs from six European radiation protection research platforms and aims to establish interaction and synergies between the different areas of expertise: MELODI (low dose and dose-rate risk research), ALLIANCE (radioecology), NERIS (emergency management), EURADOS (dosimetry issues), EURAMED (medical associations), and SHARE (social

sciences/humanities). Research findings arising from projects funded by the CONCERT calls have, along with other developments, contributed to updating the SRA. CONCERT ended in May 2020. In June 2022, the PIANOFORTE European Partnership started under the EURATOM Horizon Europe scheme and has announced yearly open calls during 2023-2025. The topics included medical exposures and risks, low dose risk more generally, inter-individual variation in response, and the effects of spatial and temporal variation in dose delivery. These areas are of direct interest to MELODI and its membership.

The activities of MELODI can be seen to be complementary to other co-ordination activities elsewhere such as the IDEA initiative in the USA, the Japanese PLANET initiative and the OECD's Nuclear Energy Agency Committee on Radiation Protection and Public Health (<https://www.oecd-neo.org/rp/crpph.html>) which established a High Level Group on low dose research (HLG-LDR) to co-ordinate efforts on a global scale, with emphasis on radiation and chemical Adverse Outcome Pathway (AOP) development, communication strategies on low dose risk research, and the development of a low dose research projects register/database. In 2022, the US National Academies of Sciences, Engineering and Medicine published a report calling for the revitalisation of low dose radiation research in the USA. This call has since been followed up by the Department of Energy, which launched the Integrated Biological and Computational Low-Dose Radiation Research programme in 2024, representing a major new initiative in this field.

The current 12th MELODI SRA for the year 2026 describes two research topics and two cross-cutting topics (which are relevant for both of the research topics) in low dose or low dose-rate radiation risk research. The topics relate to the health outcomes of concern: (1) cancers and (2) non-cancer diseases. The cross-cutting topics that are relevant to both of these disease categories are (3) individual variation in risk and (4) effects of spatial and temporal-variation in dose delivery on disease risk. Each of these is considered in detail in the SRA.

The research required to improve the evidence base for each of the four topics may be grouped into two categories:

- 1) Research to improve understanding of the mechanisms contributing to radiogenic diseases following low dose and dose-rate exposures
- 2) Epidemiological research that integrates, where possible and informative, biological and molecular markers to improve health risk evaluation of radiation exposure

The current and former versions of the MELODI SRAs and statements can be downloaded from the website: www.melodi-online.eu. The current SRA structure is outlined in **Figure 1**.




Figure 1: Outline of the structure of the Strategic Research Agenda (SRA). Numbers in parentheses refer to the SRA section dealing with each topic/issue.

2. Importance of low dose radiation health risk research

Exposure to ionizing radiation is unavoidable. Everybody encounters exposure from a range of natural and artificial sources. Medical and natural sources are the largest components of the average dose received by the general public. Exposures to artificial or natural sources can vary between individuals depending on their occupation (e.g. employment in the nuclear industry, in air transport and in medicine, particularly interventional radiologists), medical exposures (diagnostic and therapeutic procedures) and in some cases due to environmental contamination. Exposure from naturally occurring radiation involves background γ -radiation from terrestrial and cosmic sources, and internal exposure from radioisotopes such as radon and uranium (there is notable geographic variation in radon exposure). There are many and varied uses of radiation in modern society. Nuclear power generation is viewed as a carbon neutral and efficient energy source; industrial radiography plays important roles in safety assessment; medical uses of radiation for diagnostics and therapy are extensive and rapidly increasing. Long distance air travel can lead to exposures, typically 0.08 mSv for a transatlantic flight though altitude, duration and other parameters, including latitude, can affect the actual exposure level. Other sources are exposures to 'NORM' (naturally occurring radioactive materials) in the oil extraction and other industries. Broadening access to space travel is anticipated, with longer exploratory missions likely, and commercial space travel that has recently commenced.

Not only is exposure to ionizing radiation unavoidable and variable in the population, but it is known to damage health at certain exposure levels. At very high doses, radiation exposure can be lethal, while tissue damage can also occur following more localized high-dose exposures. Whole-body exposures at such levels are very rare, but for localised exposures, severe tissue damage can be observed in some patients following radiotherapy for cancer.

Evidence accumulated over many decades demonstrates that radiation can cause cancer in humans following acute exposure in the dose range of a few Gy down to 100 mGy or less, with children often showing higher sensitivity. There are indications that these more moderate exposures may also be associated with other conditions such as circulatory diseases, cataracts, cognitive impairment, immunological effects – collectively described as 'non-cancer diseases' – and possibly effects on future generations (hereditary or transgenerational effects). The risks to humans are established down to around 100 mGy for cancer in adults, about 500 mGy for circulatory diseases and lens opacities, and about 200 mGy for defects in brain development and cognition after prenatal exposure during neurogenesis. The risks to human health below these levels, especially following protracted or other non-homogenous exposures are less certain. Nevertheless, large epidemiological studies (EPI-CT) of children and young adults exposed to CT scans show statistically significant increased risk of haematological malignancies at mean doses as low as 8 mGy. Currently, the system of radiation protection aims to avoid tissue injury and minimize the risk of cancer and the possibility of hereditary disease. For radiation protection purposes, risks of cancer and possible hereditary effects below 100 mGy are regulated on the basis of an assumed linear non-threshold (LNT) relationship between dose and incidence. However, there remains uncertainty about the exact dose-response relationship for such low-dose exposures, and the impact of protracting exposures over

long periods such as during a working lifetime, and vulnerabilities of specific groups that may be at higher risk.

Striking the appropriate and acceptable balance between the benefits accrued from activities involving exposure to radiation on the one hand and the health risks posed on the other is important. The regulation for protection of individuals and populations comes at a financial cost – there are, therefore, disadvantages to both under- and over-protection. This applies in all situations – existing elevated exposure situations such as high radon areas, occupational settings such as the nuclear industry and the medical sector, and accidental situations where difficult decisions on the implementation of protective actions such as sheltering and evacuation are required. In all these contexts, it is critical to utilise robust and accurate information on the magnitude of health risks posed by given radiation doses, ranging from high to low.

The main uncertainties in radiation health risk evaluation are in the magnitude of cancer risk at low and protracted doses below 100 mGy, the magnitude of non-cancer effects below 500 mGy, the variation in individual risk within the population, and the variation in risk with dose distribution in space and time. These are therefore the key areas requiring further exploration to provide better and more reliable evidence for appropriate decision making in all areas of radiation protection. Accurate and reliable human health risk estimation at low doses is an essential foundation for a robust and acceptable system of radiation protection.

2.1 Dose and dose rate ranges to be considered

For the purposes of this document, MELODI considers low doses to be those where there remains substantial uncertainty on the magnitude of health risk. For low-LET radiations, these are taken to be those of 100 mGy and below, when considering cancer risks, and 500 mGy and below when considering non-cancer diseases as recognised by international organisations such as the United Nations Scientific Committee on the Effects of Atomic Radiation (UNSCEAR) and the International Commission on Radiological Protection (ICRP). In the context of cancer risk, moderate doses are those between 100 mGy and 2 Gy, with high doses being those above 2 Gy. For non-cancer diseases, MELODI considers moderate doses as those between 500 mGy and 5 Gy, and high doses those above 5 Gy. Low dose-rates are those of 0.1 mGy min^{-1} or less for low-LET radiation, or one-track traversal per cell per hour for high-LET radiations. These definitions apply to both organ and whole-body doses. These dose ranges are similar to many of those of interest in the medical community as set out in EURAMED documents such as the evolving research agenda under development in the EURAMED Rocc-N-Roll project (see [here](#)) and <https://roccnroll.euramed.eu>). Note that units of Sievert (effective dose, a radiation quality and tissue sensitivity weighted quantity) are frequently used for quantification of cancer risk. Effective dose, as defined by ICRP, relates specifically to cancer and hereditary effects; it is therefore not appropriate to use for non-cancer outcomes. Sieverts are also not directly measurable and so the absorbed dose units of Gray are generally used in this document; furthermore, these are the units used for dose quantification in experimental and epidemiological investigations.

3. MELODI

The purpose of the MELODI Association, as given in its Statutes, is to constitute a European Research Platform in the field of health risk assessment of low-dose ionizing radiation and its application for radiation protection as well as to coordinate and promote research and long-term competence on effects and risks to human health associated with low-dose and low-dose rate exposures to ionizing radiation.

MELODI currently has 37 institutional members including national bodies responsible for defining, funding and implementing research on low-dose risk, as well as universities and research institutes committed to contribute to R&D efforts. It is a research association that contributes to the definition of priority objectives in low-dose risk research, fostering of research programmes and initiatives to achieve these objectives, assessment of results obtained, and promotion of communication on these issues between the various parties involved, as well as ensuring sustainability of key research capacity, competencies and infrastructure. These functions are performed by organizing scientific and stakeholder workshops, promoting the visibility of the research area, establishing working groups on specific topics and facilitating collaborative research.

To achieve these goals, the establishment and regular updating of a long-term (>20 years) SRA for research to improve protection from low-dose health risks in Europe remains a major activity of MELODI. It provides guidance on the priorities for national and European research programmes and the preparation of competitive calls at the European level. Furthermore, MELODI supports the availability and maintenance of key infrastructures as an essential basis for research activities, and the retention and development of competences in radiation research and health risk assessment in the long term via an integrated European approach for training and education. As the primary aim of the MELODI SRA is to provide Euratom, national authorities and funding agencies with scientific research agendas to guide the preparation of calls and areas for prioritization, the significance of this work should periodically be evaluated for its impact on the content of calls and research prioritizations, and advances made through funded projects. Ultimately, the research guided by the SRA is anticipated to make an impact on radiation protection policy.

Following the recommendations and roadmap established by the High Level and Expert Group on European Low Dose Risk Research (HLEG) in 2009 (full report [here](#); for a summary see [Repussard, 2018](#)), and supported over time by DoReMi, OPERRA, CONCERT, and PIANOFORTE, the latter moving to integrate SRAs covering all aspects of radiation protection research in Europe, the radiation protection research community within Europe has been progressively more deeply integrated over the past decade.

MELODI participates in the body representing all radiation protection platforms, MEENAS; this body serves to coordinate the entire range of radiation protection research areas in Europe. An important development has been the publication of a joint-platforms roadmap for radiation protection research.

In October 2010, the first draft of a MELODI SRA was published on the MELODI Website and opened for public consultation. The contents were based on the considerations and key priority issues formulated by the HLEG and DoReMi. In February 2014, the MELODI Board of Directors (BoD, now re-constituted as the MELODI Executive Committee) established three working groups (WGs), on the MELODI SRA,

Education & Training, and Infrastructures. The MELODI SRA is updated periodically by the SRA WG, taking into account recent and emerging research results and radiation protection research issues. The updated draft is posted on the public MELODI website, usually before the annual MELODI workshop - now European Radiation Protection Week (ERPW). An open consultation process is set up via the website and the workshop to seek input from other scientists and stakeholders before the SRA's revision. The updated SRA is also sent to the independent Scientific Advisory Committee of MELODI for comment. The final SRA is prepared for approval by the MELODI Executive Committee. The current edition of the SRA will be the twelfth version.

3.1 MELODI in the context of other radiation research platforms

Large parts of European radiation protection research have been organized within the CONCERT European Joint Programme (EJP), and the ongoing PIANOFORTE project continues to develop this coordinated approach. The EJP and PIANOFORTE European Partnership have brought, and will continue to bring, together relevant funding agencies from the EC and Member States to integrate European research, and to administer calls for research proposals in radiation protection on behalf of the European Commission. This activity builds upon the Strategic Research Agendas from six European radiation protection research platforms, MELODI, ALLIANCE (radioecology), NERIS (emergency management), EURADOS (dosimetry issues), EURAMED (medical associations), and SHARE (social sciences and humanities), and aims to establish interactions and synergies between the different areas of expertise. Integration across the different platform areas is being fostered and developed through the drafting of a PIANOFORTE Strategic Research and Innovation Agenda (SRIA) to guide all research related to radiation protection (<https://pianoforte-partnership.eu/about/sria>). This SRIA is expected to be implemented as part of the Euratom Research and Training Programme 2026-2027 prolongation of the PIANOFORTE project. MELODI also aims to be responsive to the key challenges in the system of radiation protection as identified by international organisations such as UNSCEAR and ICRP.

4. List of recent projects

The following table provides a non-exhaustive overview of recently completed and ongoing European projects relevant to radiation protection research and the MELODI SRA context. The descriptions are based on official project information, primarily CORDIS and project websites, and are not intended as an assessment of project outcomes.

Table 1. Recent and ongoing European radiation protection research projects relevant to the MELODI SRA.

Acronym	Full title	Programme / framework	Period / status	Official short description / abstract
MEDIRAD	Implications of Medical Low Dose Radiation Exposure	H2020-Euratom	1 June 2017 – 28 February 2022; closed	MEDIRAD aimed to enhance the scientific basis and clinical practice of medical radiation protection, including health effects of low-dose ionising radiation from diagnostic and therapeutic imaging and off-target effects in radiotherapy. It addressed organ dose estimation, medical-radiation epidemiology, biomarkers, mechanisms and risk models. (CORDIS)
HARMONIC	Health effects of cArdiac fluoRoscopy and MOderN radIotherapy in paediatricCs	H2020-Euratom	1 June 2019 – 30 November 2024; closed	HARMONIC investigated cancer and non-cancer outcomes after medical ionising-radiation exposure in young cancer patients undergoing external beam radiotherapy and children with cardiac defects receiving cardiac fluoroscopy procedures. (CORDIS)
RadoNorm	Towards effective radiation protection based on improved scientific evidence and social considerations — focus on radon and NORM	H2020-Euratom	1 September 2020 – 31 August 2025; closed	RadoNorm addressed protection from harmful effects of ionising radiation and radioecological impacts of radionuclide contamination, aiming to reduce key uncertainties in low-dose radiation risk assessment with a focus on radon

Acronym	Full title	Programme / framework	Period / status	Official short description / abstract
SINFONIA	Radiation risk appraisal for detrimental effects from medical exposure during management of patients with lymphoma or brain tumour	H2020-Euratom	1 September 2020 – 31 August 2024; closed	and NORM, from molecular mechanisms to human health and biota. (CORDIS) SINFONIA developed methodologies and tools for comprehensive risk appraisal of detrimental effects of radiation exposure on patients, workers, carers, the public and the environment during management of lymphoma and brain-tumour patients. It also addressed patient variability in radiation sensitivity and secondary malignancy risk. (CORDIS)
EURAMED rocc-n-roll	EUROpeAn MEDical application and Radiation prOteCtion Concept: strategic research agenda aNd ROadmap interLinking to heaLth and digitisation aspects	H2020-Euratom	1 September 2020 – 31 August 2023; closed	EURAMED rocc-n-roll analysed research and radiation protection needs in medical applications of ionising radiation, produced an SRA and roadmap for medical applications and related radiation protection, and considered synergies with existing radiation protection platforms and Euratom-funded projects. (CORDIS)
PIANOFORTE	Partnership for European research in radiation protection and detection of ionising radiation: towards a safer use and improved protection of the environment and human health	Horizon Europe / Euratom European Partnership	1 June 2022 – 31 May 2029; ongoing	PIANOFORTE is the current European partnership for radiation protection research. It supports research and innovation for improved protection of the public, patients, workers and the environment, including through open calls funding research projects. (CORDIS)
DISCOVER	Dissecting radiation effects into the Cerebellum microenvironment driving tumour promotion	PIANOFORTE-funded research project	2023 call	Uses Ptch1 ^{+/-} mice, cerebellum slices and cell cultures to study how cerebellar populations respond to moderate and low radiation doses

Acronym	Full title	Programme / framework	Period / status	Official short description / abstract
SONORA	Safe, optimized and personalized radiology and radiotherapy procedures for pregnant patients	PIANOFORTE-funded research project	2023 call	and how the microenvironment transmits radiation signals involved in carcinogenesis. Addresses medical exposure of pregnant or potentially pregnant patients, with the aim of producing guidance for fetal dose estimation in radiology and radiotherapy procedures.
LutADose	Personalized dosimetry to improve the clinical outcome of prostate cancer patients treated with $^{177}\text{Lu}/^{225}\text{Ac}$ -PSMA targeted therapies	PIANOFORTE-funded research project	2023 call	Develops image-guided dosimetry for $^{177}\text{Lu}/^{225}\text{Ac}$ -PSMA radioligand therapy so that therapeutic doses can be tailored to individual patients and improve the risk-benefit balance.
IMAGEOMICS	Optimizing Benefit/Risk Ratio in Breast Cancer Diagnosis and Radiotherapy: Identifying Molecular, Cellular and Imaging Signatures of Breast Cancer Heterogeneity to Improve Personalized Therapeutic Strategies for Synergistic Treatment Combinations	PIANOFORTE-funded research project	2023 call	Aims to improve the benefit/risk ratio for breast cancer patients by identifying response signatures for combined radiotherapy and immunotherapy and developing imaging with increased diagnostic potential and reduced ionising radiation exposure.
CORNET	deCIPHERing the rOle of micRoenvironment after low-dose exposure for coloN carcinogenEsis and radiaTion risk	PIANOFORTE-funded research project	2024 call	Investigates the role of the microenvironment after low-dose exposure in colon carcinogenesis and radiation risk.
UBT-Rad	Unraveling brain tumor formation after low dose irradiation exposure	PIANOFORTE-funded research project	2024 call	Studies mechanisms underlying low-dose-ionising-radiation-induced brain carcinogenesis, including the role of the brain microenvironment and early biomarkers.

Acronym	Full title	Programme / framework	Period / status	Official short description / abstract
MIRAMARE	Mechanisms of the inversed relationship between menarche age and radiation-induced breast and endometrial cancer	PIANOFORTE-funded research project	2024 call	Studies radiation-induced breast and endometrial cancer in relation to modifiers such as menarche age, obesity and hormones, including acute and chronic exposure for low-dose-rate risk assessment.
KAYAC+	Knowledge on outcome of adolescents and young adults with cancer	PIANOFORTE-funded research project	2024 call	Collects knowledge on outcomes of adolescents and young adults with cancer; the official project information notes cancer incidence in AYAs and treatment/follow-up data needs.
DOSELIA	Computing whole-body radiation dose distributions and subsequent cancer risks from modern radiotherapy techniques in paediatric patients	PIANOFORTE-funded research project	2024 call	Aims to improve assessment of whole-body doses and subsequent primary cancer risks in paediatric patients undergoing external beam radiotherapy, including out-of-field doses and paediatric risk projection models.
EMPATHY	Evaluation and optimization of proton arc therapy	PIANOFORTE-funded research project	2024 call	Addresses optimisation of proton arc therapy and adaptive proton therapy, including challenges from low-dose dispersion over larger tissue volumes and more intensive X-ray imaging.
BIOMBRT	Unravelling the biological mechanisms of normal tissue preservation in spatially fractionated minibeam radiation therapy	PIANOFORTE-funded research project	2025 call	Investigates biological mechanisms of normal-tissue preservation in minibeam radiation therapy, including effects in radiosensitive organs such as brain and lung.
FORRCE	Framework for outcomes of radionuclide radiation on cellular environment	PIANOFORTE-funded research project	2025 call	Develops and validates a mechanistic framework for targeted radionuclide therapy, linking energy deposition with biological outcomes across spatial and temporal scales,

Acronym	Full title	Programme / framework	Period / status	Official short description / abstract
PRESTO	PRoton therapy Enhancement by Spatially fractionated Treatment Optimization	PIANOFORTE-funded research project	2025 call	including low dose rates and heterogeneous energy deposition. Investigates proton minibeam radiation therapy, aiming to maintain tumour control while preserving spatial fractionation patterns that protect normal tissues.

Additionally, MELODI has sponsored eight topical workshops, concerning (i) individual radiosensitivity and radiosusceptibility, (ii) non-cancer effects of ionising radiation, (iii) the effects of spatial and temporal variation in dose delivery, (iv) adverse outcome pathways for radiation effects, (v) effects of ionising radiation exposure in offspring and next generations, (vi) radiation-induced circulatory diseases, (vii) radiation effects in the central nervous system, and (viii) radiation effects on the immune system. They are summarized in Table 2. Outputs and recommendations from several workshops have been published. The full bibliographic records can be found in the Annex. Further publications from the more recent workshops are anticipated. Key recommendations from the published outputs have been taken into consideration in developing this SRA.

Table 2. MELODI-sponsored topical workshops and related publication outputs.

Title	Location	Date	Organised by	Publications
MELODI Workshop on Individual Radiosensitivity	Malta	12-14 March 2018	CEA	Salomaa and Jung (2020); Gomolka et al. (2020); Averbeck et al. (2020); Seibold et al. (2020); Kalman and Oughton (2020)
MELODI Workshop on non-cancer effects of ionising radiation	Sitges, Spain	10-12 April 2019	ISGlobal	Kreuzer and Bouffler (2021); Ainsbury et al. (2021); Lumniczky et al. (2021); Pasqual et al. (2021); Tapio et al. (2021)

Title	Location	Date	Organised by	Publications
MELODI Virtual Workshop on the Effects of Spatial and Temporal Variation in Dose Delivery	Virtual	17-20 November 2020	Centre for Energy Research	Madas and Wojcik (2022); Pazzaglia et al. (2022); Lowe et al. (2022); Baiocco et al. (2022); Madas et al. (2022); Li et al. (2022); Saldarriaga Vargas et al. (2024)
MELODI Workshop on Adverse Outcome Pathways for radiation effects	Virtual	13-15 April 2021	IRSN, jointly with ALLIANCE and OECD HLG-LDR	Chauhan et al. (2022a); Chauhan et al. (2022b); Chauhan et al. (2022c); Chauhan et al. (2022d); Azimzadeh et al. (2022); Jaylet et al. (2022); Burt et al. (2022); Kozbenko et al. (2022); Yu et al. (2022); Klovov et al. (2022); Tollefsen et al. (2022); Stainforth et al. (2022); Adam et al. (2022); Chauhan et al. (2021)
Effects of Ionising Radiation Exposure in Offspring and Next Generations	Budapest	31 May-2 June 2022	Sisko Salomaa; jointly with ICRP and ALLIANCE	Streffer and Hande (2024); Amrenova et al. (2024a); Amrenova et al. (2024b); Benotmane and Trott (2024); Degenhardt et al. (2024a); Grison et al. (2024); Nakamura et al. (2024); Sreetharan et al. (2024); Stephens et al. (2024); Liutsko et al. (2024); Zölzer et al. (2024); Degenhardt et al. (2024b)
Updates on Radiation-Induced Circulatory Diseases	Virtual	30 May-2 June 2023	BfS	Hamada (2023)
Radiation Effects in the Central Nervous System	Rome	18-19 February 2025	ENEA	Nakamizo (2025); Hamada et al. (2026); further review publications under preparation
Radiation Effects on the Immune System	Grenoble, France	28-29 April 2026	CEA	review publications under preparation

Beyond the activities of MELODI, documents of relevance to this SRA have been published by UNSCEAR in recent years – most notably UNSCEAR 2019, [Annex A](#) - Evaluation of selected health effects and inference of risk due to radiation exposure and [Annex B](#) - Lung cancer from exposure to radon and the 2020/21 [Annex C](#), Biological mechanisms relevant to the inference of cancer risks from low dose and low dose-rate radiation. In 2021, ICRP published a paper outlining its approach to review of the system of radiological protection ([Clement et al, 2021](#)) and in the same year published a paper summarising the areas of research that are expected to support the development of the system of protection ([Laurier et al, 2021](#)). NCRP published in 2020 its Report No. 186 – Approaches for Integrating Information from Radiation Biology and Epidemiology to Enhance Low-Dose Health Risk Assessment (2020) that also places emphasis on AOPs and related approaches.

5. Strategic Research Agenda

Radiation protection is one particular area of health protection concerning the prevention of radiation-induced diseases and tissue damage, notably cancers and some non-cancer diseases in the general public, patients and workers. The health impacts of radiation generally concern diseases or biological effects that are multi-factorial in origin, with both intrinsic and extrinsic risk factors. The intrinsic, non-modifiable risk factors include age and sex, as well as less well characterised genetic and other factors, all of which, in addition to being important risk factors in themselves, may also modify the health impact of radiation. There is also a wide range of modifiable risk factors affecting the incidence of these diseases, including behavioural factors such as diet, tobacco smoking and exercise, as well as natural and human-made environmental factors including co-exposures to other environmental and occupational carcinogens and medicinal drugs. Radiation protection research therefore needs to be viewed in this wider context where any radiation exposures and effects on health rarely, if ever, occur independently; rather individuals and their disease risk can be influenced by their genome, epigenome, exposome, microbiome and other factors. This wide range of influences on individual and population health risks can pose problems for discerning the impact of radiation exposures, especially when exposure levels are low. As stated earlier, radiation protection is one element of general health protection relevant in public, occupational and medical exposure settings.

The MELODI SRA is based on the key policy goals defined by the High Level and Expert Group to address the robustness of the current radiation protection system with evolution to take account of developments in the area (see **Figure 2**).




Figure 2: Key policy issues in European low dose radiation risk research defined by the High Level and Expert Group.

The key policy issues identified in the HLEG report are:

- The shape of the dose-response for cancer;
- Tissue sensitivities for cancer induction;
- Individual variability in cancer risk;
- The effects of radiation quality (type);
- Risks from internal radiation exposure;
- Risks of, and dose response relationships for, non-cancer diseases and hereditary effects.

For the purpose of the MELODI SRA, these issues were restructured into two topics relating to disease types and two cross-cutting issues (**Figure 1**):

TOPICS

- (1) Dose and dose-rate dependence of cancer risk;
- (2) Risk for non-cancer effects;

CROSS-CUTTING ISSUES

- (3) Individual variation in risk;
- (4) Effects of spatial and temporal-variation in dose delivery

As discussed by the HLEG and confirmed by MELODI, research at low dose-rates or low doses presents significant challenges in the investigation of both radiation-related health effects and underlying biological mechanisms, because both the magnitude of health risk and the biological effects are expected to be small. A multidisciplinary approach is therefore essential. Both epidemiological and experimental studies will also require

sufficient statistical power and sample sizes to be effective. Nested case-control study designs are likely to be suitable, as well as “meet in the middle approaches” for example.

For these reasons, discussion of each key question is sub-divided below into two categories:

- Research to improve understanding of the mechanisms contributing to radiogenic diseases following low dose and dose-rate exposures.
- Epidemiological research that integrates, where possible and informative, biological approaches to improve health risk evaluation.

5.1 Dose and dose-rate dependence of cancer risk

Current risk estimates used in radiation protection are based upon epidemiological studies of exposed populations. Radiation protection standards aim to avoid tissue reactions and minimize the incidence of the late stochastic effects of cancers and possible hereditary effects in future generations. Thus, it is of fundamental importance to radiological protection that the health risk estimates are evidence-based, robust, and credible. Most important among the epidemiological studies are the follow-up studies of Japanese populations exposed as a consequence of the atomic bombings of Hiroshima and Nagasaki that provide risk estimates per unit dose for cancer and, more recently, non-cancer effects. While the Japanese studies remain the main basis for the cancer risk estimates used in radiation protection, they relate to a specific population and a specific exposure scenario. The exposure was essentially instantaneous, high dose rate, total body gamma ray exposure with a small neutron component. Information about cancer risk from the A-bomb survivor studies is, to an increasing extent, complemented by studies of occupational, environmental and medical exposures, which allow direct investigation of effects of fractionated or protracted exposures in current European populations.

As noted earlier, epidemiological studies provide direct evidence of dose-related increases in cancer risk after acute exposures with doses of about 100 mGy and above. Recent studies have provided better evidence of risk at doses below 100 mGy and with protracted exposure. Some reports indicate a possible increased risk of childhood leukaemia, and possibly brain tumours from doses below 100 mGy due to natural background gamma radiation and from paediatric CT imaging. The most recent studies on haematological malignancies suggest increased risks in (1) young people having received at least one CT scan before 22 years of age at mean doses as low as 8 mGy (a total dose of 100 mGy multiplied the risk by about a factor of 3), and (2) nuclear workers (International Nuclear Workers Study, INWORKS) having been exposed to protracted low doses with an average cumulative red-bone-marrow dose of 16 mGy.

Nevertheless, major uncertainties remain concerning (i) the magnitude of total and organ-specific cancer risks following specific exposure situations such as protracted exposure encountered in environmental, as well as occupational and medical settings, and when the dose is inhomogeneously distributed, particularly after internal contamination; (ii) the risk for individual cancer sites due to different tissue sensitivities, and (iii) the best evidence-based models to infer risk at doses and dose-rates that are lower than those for which solid

epidemiological evidence is available. In this context, there are also a number of ethical questions that need to be addressed, such as the use of the LNT model for extrapolation to very low doses, and whether other risk factors may substantially modify radiation risks.

Classical epidemiological studies will need to be continued to refine the knowledge of risk directly in human populations, particularly in the context of low dose, protracted and non-uniform exposures. Accuracy of risk estimates can potentially be increased by more precise dose estimation and outcome assessment, larger studies or pooling of data from several studies. Mechanistically, studies should investigate the contribution of epigenetic and bystander / inflammatory signalling processes to cancer development, providing new data informing on the role(s) of radiation in the different stages of radiation carcinogenesis. Radiation-induced ageing and senescence-associated inflammation might contribute to cancer risk. Mechanistic and epidemiological approaches should be integrated whenever feasible to address cancer risks from acute whole-body exposures with low-dose (<100 mGy) or from fractionated, protracted and inhomogeneous exposures resulting in low-to-moderate doses. Studies also need to address the impact of different radiation qualities and effects of both internal and external exposures, alone and in combination. Knowledge of health risks from low dose-rate exposures is of direct relevance for radiological protection in emergency situations, in medicine (with children, who are known to be more sensitive than adults, easily receiving doses of several tens of mGy to the brain from multiple brain CT scans), and in occupational settings, with the current dose limit of 20 mSv/year averaged over 5 years with no single year exceeding 50 mSv. Radiation protection in the medical context is particularly important as exposures have been increasing, and novel radiation modalities are both in use and under development. Clear and coherent principles for justifying the long-term risks with more immediate benefits are needed. Radiation protection in the context of long-term space travel, where radiation exposures have some very specific characteristics and differ from terrestrial exposures, is likely to grow in importance in the future. In the context of the risks associated with space travel, the formation of links to other relevant agencies with research programmes, e.g. NASA, ESA, JAXA, would be beneficial. Beyond studies of specific irradiated populations, major cohorts have been established and followed primarily for reasons other than the assessment of radiation risk. Some consideration of the benefits of utilising such cohorts and adding radiation exposure information may be of use in the future.

Research line: Health risk evaluation

Quantification of cancer risk at moderate dose or from acute or protracted non-uniform exposure, and at low doses or dose rates from acute homogenous exposure are key challenges for improved radiation risk assessment. The large size of epidemiological studies required to detect small increases in cancer risk at low doses and dose rates, the need to capture all major sources of radiation exposure, and the potential for bias and confounding factors present practical challenges, particularly at the lowest doses. The priorities in this area include the maintenance and improvement of key cohorts by continued follow-up, pooling of different studies, collection of information on confounders and improving precision of dose and outcome data. Key cohorts are characterized by large populations with exposure conditions and dose distributions that are relevant for radiation protection, good individual dosimetry, long and complete follow-up with good quality of health outcome data, particularly in relation to cancer occurrence; and

the possibility of collecting information on relevant potential confounders either on the whole cohort or through nested case-control studies.

These studies should include, where possible and likely to be informative, the collection and appropriate storage of a large number of relevant biological samples, including tissue samples from cancer cases and somatic tissue from affected individuals; while this is generally difficult in large-scale cohorts, it can be integrated in nested case-control studies. Through identification, validation and integration of relevant biological endpoints and markers into epidemiological studies, further insights will be gained into the risks associated with such exposures at the population and individual levels. The integration of both epidemiological and mechanistic studies will improve cancer risk evaluation through molecular epidemiological studies or by mechanistic modelling.

Priority research areas are:

- To determine the shape of the dose and dose-rate response relationships in humans for all cancers, and where possible specific cancer sites, based on key informative epidemiological studies, including medical and occupational cohorts, as well as those accidentally exposed.
- To determine the risk for different cancer types based on key cohorts (see above) in order to investigate differences in tissue sensitivity.
- To carry out studies to fully understand the cell autonomous and indirect mechanistic processes by which radiation contributes to cancer effects, since the same dose may not infer the same consequences if present in another cellular, co-exposure, inflammatory and/or genetic and epigenetic context
- To evaluate the dose-response for tumour types, ideally defined by molecular characterisation.
- To investigate precursor lesions of cancers in any available biological samples, e.g. tissue or saliva/blood and by imaging methods in study populations with well-characterised exposure to allow modelling of carcinogenesis, including AOP approaches.
- To identify and validate biomarkers of exposure and health effects related to cancer, both working from early exposure biomarkers through intermediate steps to disease as in cohort studies, and tracing markers of susceptibility, pre-clinical changes and biological indicators of exposure in disease as in case-control studies – the ‘meet in the middle’ approach.
- To determine the value of evaluating cancer risks through systems biology analyses and models of carcinogenesis based on mechanistic studies and epidemiological data, and integration of the two.

Research line: Basic mechanisms

An LNT extrapolation model is currently used to estimate risk at low doses from higher dose epidemiological data. An important aspect of the justification of using this model is that radiation carcinogenesis is assumed to be primarily driven by damage to DNA and subsequent mutation of growth-regulating genes in target cells. Mutational signatures of

radiation in tumours obtained by deep sequencing have shown potential driver as well as passenger mutations. A higher proportion of deletions was suggested as an ionising radiation-induced signature, indicating DNA damage and loss of genetic material through erroneous repair. In addition, a number of other potential mechanisms contributing to and modulating radiation carcinogenesis have been proposed, including epigenetic mechanisms of gene regulation such as, but not limited to, DNA methylation, chromatin conformation and miRNA expression, transmissible genomic instability, bystander effects, inflammation and adaptive response, pre-existing mutations, and it is of utmost importance to determine their roles. The extent to which these modulating effects and non-mutational mechanisms challenge the validity of the LNT risk extrapolation model needs to be determined. For this purpose, the use of well validated animal and human cellular / tissue models of radiation carcinogenesis (both solid cancers and leukaemias) is required.

Priority research areas are:

- To carry out studies to fully understand the cell autonomous and non-targeted mechanistic processes by which radiation contributes to cancer effects, since the same dose may not infer the same consequences if present in another cellular, co-exposure, inflammatory or genetic and epigenetic context.
- To determine the nature, roles and radiosensitivity of the various target cells for radiation carcinogenesis. The most important of these are generally taken to be stem and progenitor cell populations, which may have specific responses to radiation.
- To determine the contribution of DNA damage / mutational processes at low doses and dose-rates and with differing radiation qualities. Further information on the specific genes affected at low doses in the development of specific cancers and quantitative aspects can contribute to refining risk extrapolation models and the identification of radiation exposure and cancer biomarkers.
- To determine the contribution of epigenetic modifications and cellular ageing, including senescence-associated inflammation. Gene function and cellular processes can be regulated at the epigenetic level. The extent to which radiation affects epigenetic states that relate to carcinogenesis needs to be elucidated, and also how epigenetic factors affect response to radiation.
- To determine the influence of cell micro-environmental, non-targeted and systemic processes that may promote or restrict the growth of pre-malignant cells in tissue, and how radiation exposure affects the tissue environment to facilitate or retard the growth of (pre)-malignant cells. For example, the influences of low dose radiation exposure on inflammatory reactions, which may promote a cancer-prone environment, and effects of radiation on immune-cell homeostasis and function, which may impact immune surveillance against cancer cells.
- To examine the extent to which any of the above are different at high dose / dose-rate by comparison with low dose / dose-rate. All these investigations are likely to benefit from the application of Artificial Intelligence/Machine Learning (AI/ML) approaches to analyses of multi-omics and big data sets in the future.

5.2 Non-cancer effects

It has been traditionally assumed that health effects other than cancer and hereditary diseases show a threshold (defined as the dose required to lead to 1% excess incidence) at doses that are well above the levels of exposure typically encountered in the public environment, at work or in diagnostic medical uses of ionizing radiation. Recent results from epidemiological and experimental studies indicate possible increased risks of circulatory diseases, cataracts, cognitive/neurological effects and others not only at high doses but also down to 500 mGy and, possibly even lower. Based on these findings the ICRP issued in 2011 a statement on tissue reactions (formerly termed non-stochastic or deterministic effects) that noted evidence that the threshold in absorbed dose for effects on the lens of the eyes is of the order of 500 mGy (acute and protracted exposure). Consequently, a recommendation was made for a reduction in the annual equivalent dose limit for the eye lens to 20 mSv per year averaged over 5 years, with no one year exceeding 50 mSv. In addition, ICRP suggested that the dose threshold for circulatory diseases may be as low as 500 mGy.

Evidence for radiation-related hereditary effects is based on experimental animal studies. There is no direct evidence for hereditary/transgenerational effects from human studies, though 2nd and 3rd generation studies are likely to be feasible in specific cohorts, e.g. in the Urals. The as yet uncertain contribution of hereditary risk to overall risk is expected to be small in comparison with that of cancers. While the system of radiation protection includes hereditary effects in the calculation of low dose detriment along with cancers for risk management purposes, the range of diseases occurring among the offspring of irradiated parents includes both cancer and non-cancer diseases. There are also effects on pregnancy outcomes, and as such the pregnant mother represents a high risk group.

For all outcomes, there are uncertainties about possible effects at low doses, which may have important implications for radiation protection. Results of available epidemiological studies are not always consistent, as the risk estimates are prone to bias and confounding, and the biological mechanisms of relevance for health risks at these low doses are not fully understood. The possibility of a stochastic nature of non-cancer effects without dose thresholds raises a wide range of questions and needs further investigation. In contrast to cancer, knowledge on the underlying biological mechanisms for radiation-related non-cancer effects in the moderate and low dose range is sparse. Therefore, research to understand the mechanisms is necessary. In addition, epidemiological research of key cohorts with good information on potential confounding factors is needed to provide information on radiation-related risk of non-cancer diseases following low-dose, protracted or fractionated exposure, relevant for radiation protection. Individual variation in risk, mixed exposures and impact of characteristics of radiation exposure will also need to be explored.

Research line: Health risk evaluation

Quantification of non-cancer disease risk in humans at moderate or low doses or dose-rates is a key and difficult challenge for radiation protection, because the magnitude of risk due to radiation is expected to be low and the potential for bias and confounding factors is high. Informative epidemiological studies in this field will be characterized by cohorts of large size with exposure settings and dose levels relevant for radiation protection, good dosimetry, high quality of health data, long follow-up and the possibility of collecting

information on relevant potential confounders either on the whole cohort or through nested case-control studies. In addition, these studies should include – where possible and informative – collection of biological samples, relevant tissue samples from the relevant organ to allow mechanistic investigations, and extensive data on the health status during follow-up.

Through improvement of key epidemiological studies (e.g., increasing the statistical power by pooling studies using standardized study protocols; development of new strategies for data acquisition (automated data collection, data from health insurance providers); improvement of appropriate organ and tissue dose assessment, e.g. different parts of the heart, main arteries and veins, as well as blood, brain, eye lens, etc.) and, where possible and informative, the identification and integration of relevant biological endpoints and markers into epidemiological investigations, further mechanistically-informed insights will be gained into the risks associated with such exposures. The creation of a regularly updated database on epidemiological studies of low-dose radiation effects on non-cancer outcomes could significantly enhance low-dose radiobiology. It would facilitate data access, support meta-analyses, improve risk assessment, and contribute to better radiation protection guidelines.

Priority research areas are:

- To determine the shape of the dose-rate and dose-response relationships, notably the presence or absence of threshold doses (down to a few tens of mGy), in humans for non-cancer outcomes at low or moderate doses based on key informative epidemiological studies (molecular or otherwise, as appropriate). While increasing numbers of studies concern circulatory diseases, few are available on cognitive impairment, neurodegeneration and neuropathies as well as on hereditary/transgenerational effects. Any such studies require careful and explicit definition of the disease outcomes being assessed. Because of the long lag times there is a need for more comprehensive, long-term investigations and/or the development of novel biological models to facilitate the study of these outcomes within shorter timeframes.
- To identify, develop and validate biomarkers of exposure (especially for low doses and protracted/inhomogeneous exposures), and for early and late non-cancer effects. Relevant tissue banks are currently available but need to be further expanded to allow for investigation of more possible exposure scenarios that can lead to low-dose-induced non-cancer effects. The development of such biomarkers would allow better estimation of the actual doses received and inform the evaluation of the dose-response relationship of non-cancer effects.
- To investigate early stages in the progression of non-cancer effects in tissue or disease-related endpoints in biological samples from members of appropriate epidemiological studies or individuals with similar living conditions and known exposure in order to understand spontaneous pathogenesis. This is a pre-requisite to understand radiation effects on pathogenesis.
- To evaluate non-cancer risk through systems biology analyses and mathematical models combining and integrating mechanistic studies and the epidemiological data.

Research line: Basic mechanisms

Deterministic effects or tissue reactions are classically thought to arise as a consequence of cell killing or functional inactivation by high radiation doses. They are characterised by steeply increasing dose-response relationships at doses exceeding a defined threshold. It is unlikely that cell killing/inactivation will explain fully the effects of lower radiation doses on circulatory diseases, cataract and cognitive dysfunction. Low dose radiation may induce cellular senescence, and it may be debatable if this consequence of DNA damage also could be regarded as a stochastic event. The occurrence of this phenomenon in tissue stem cell compartments is an event that could have profound pathophysiological consequences. Alteration of stem cell functions may impair tissue renewal and homeostasis or may promote non-cancer diseases or cancers. Epidemiological investigations require support from studies to identify the underlying mechanisms that lead to each of the non-cancer diseases. Each disease may have a different mechanistic basis, and it is not clear whether there will be any similarity with the mechanisms that lead to radiation-related cancers.

Priority research areas are:

- To integrate classical in vitro and in vivo biological models with innovative experimental approaches—such as organoids, assembloids, organ-on-chip, 3D bioprinted models, and in silico integrations—that more accurately mimic human physiology. This integration should bridge the gaps between traditional in vitro and in vivo models, as well as between non-human in vivo models and humans. Emphasis should be placed on developing models for target tissues affected by radiation-related non-cancer diseases (e.g., circulatory diseases, cataracts, neurological dysfunctions) to elucidate underlying pathways. Additionally, these models should provide insights into mechanistic differences between acute and chronic exposures and varying radiation qualities. The ultimate goal is to enhance collaboration and complementarity in radiation research by combining conventional models with advanced 3D human cell-based approaches.
 - To investigate the early stages of radiation-induced non-cancer diseases, aiming to link molecular and cellular effects to impairments at the organ level. AOP concepts for the spontaneous development of these diseases, as well as radiation-related AOPs (expanded with quantitative data) should be included. This knowledge will enable the identification of bioindicators for pathological processes, which can subsequently be validated in molecular epidemiological studies.
 - To investigate the effects of low-dose radiation on human health, considering exposure to mixed fields (i.e. space radiation, radon exposure, nuclear accidents, nuclear medical applications, etc.).
- To apply a full range of analytical methods including multi-omics and multi-modal data integration (with application of AI/ML where appropriate) and consideration of the target cells and surrounding microenvironment. In this context, emerging technological innovations including single cell multi-omics may help to identify differences in radiation sensitivity between relevant cells and tissues. The mechanistic knowledge gained is likely to be useful for the identification of relevant biomarkers,

e.g. specific metabolic and pathological changes that are clearly radiation-induced, and the development of mechanistic models of disease development.

- To investigate the role of low-dose and low dose-rate radiation-induced senescence in neuronal stem and progenitor cells as a driver of neurodegeneration and cognitive decline, including the underlying molecular pathways, the impact on neurogenesis, radiation effects on supporting cells, and the potential contribution of the senescence-associated secretory phenotype (SASP) to chronic neuroinflammation and neuronal dysfunction.
- To determine if other pre-existing conditions, such as neuropathies, inflammatory conditions or metabolic and mitochondrial diseases, but also non-modifiable factors (such as sex, age and genetic background) affect the incidence of radiation-induced non-cancer outcomes

5.3 Individual variation in risk

Individual variation in radiation-related cancer and non-cancer disease risk is a key area to address for radiation protection; effective identification of groups at higher risk is necessary to ensure that adequate protection is provided to all. Differences in the magnitude of radiation-induced risks between individuals, or groups, may relate to sex, age at exposure and age attained, state of health, genetic (including both germline and somatic) and epigenetic make-up, as well as behavioural factors. Such differences, if significant, raise very important ethical and policy questions as to whether some individuals or groups are or will be inadequately or over-protected by the present system and regulations. Similar concerns on variation in risk between individuals apply to non-cancer outcomes. These ethical considerations might, for example, be important in the context of space travel, where individual-level risk estimates could potentially be considered in crew selection.

At present, there is insufficient information about the magnitude of the differences in radiation response between individuals or groups of individuals and their consequent influence on risk estimates at low doses and dose-rates. In order to address policy questions, it is necessary to obtain better scientific information on the extent of the variations in radiation response (sensitivity to radiation including susceptibility) in the population, in the magnitudes of the variations, characteristics affecting the variation and in the sizes of the various population subgroups. Importantly, reliable and robust biomarkers predictive of individual risk need to be identified and characterised through basic mechanistic research before application in epidemiological studies.

The group of individuals who reach a high age is expanding in many countries worldwide. This may increase the need for additional considerations when estimating long-term risks after radiation exposure; this is particularly true in light of new findings on the existence of pre-leukaemic mutations in a relatively large proportion of older individuals for example. Further research on additional factors such as immune system alterations responsible for expansion of mutated clones aims to improve understanding of risks of exposure.

Diagnostic procedures with low-dose radiation are increasing worldwide. These procedures are beneficial to the patient if properly justified. Nevertheless, there is scant awareness of health risk associated with their unjustified or excessive use. For instance,

health risk may be higher in elderly patients due to increased vulnerability, the possible impact of co-morbidities and poor recovery of homeostasis following a stress such as low dose radiation exposure. In certain scenarios and tissues however, risks may be highest when children or adolescents are exposed, in particular during time periods when cell proliferation within a particular tissue is rapid.

Consideration of how individual differences affect the relationship between absorbed dose (and dose distribution) and risk is required. For internal intakes of radionuclides, the dose and dose distributions can be very different in individuals for the same exposure because of anatomical and physiological differences (e.g. in airway morphology or breathing mode). These variabilities should be taken into account by accurate dosimetric and physiologically relevant biokinetic models. In addition, the nature of the interaction of ionizing radiation with co-exposures to other agents (e.g. tobacco smoke, heavy metals and, generally, environmental pollutants) and existing risk-modifying conditions (e.g. iodine deficiency for thyroid cancer) for the onset of various cancers and diseases are important in considering risk transfer between different populations.

Research line: Health risk evaluation

The quantification of the contribution that individual variation in response to radiation makes to radiation health risk on both an individual and population level is a key question. Realistic estimates of the magnitude of differences in response between individuals and groups are needed.

Priority research areas are:

- To identify and validate candidate biomarkers of individual response identified from mechanistic or clinical studies in cohorts of exposed and non-exposed subjects who have developed cancers or non-cancer diseases. Combinations of biomarkers may present the highest stringency. As few suitable large cohorts with biological samples are currently available, proof-of-principle studies with cohorts exposed to higher doses should be conducted to refine methodologies and to extrapolate to low doses. Links to existing biobank resources, particularly in relation to medically exposed patients should be considered.
- To improve or set up molecular epidemiological cohorts or case-control studies to determine factors (host and environmental) that modify individual risk of radiation-induced cancer and non-cancer effects and quantify their impact.
- To quantify the variation in risk between different population groups and the impact of different factors, for example, age at exposure and attained age, as well as co-exposures and host factors, including anatomical, physiological and cell type differences. Knowledge of the nature of possible interactions between ionizing radiation and these factors on health risk (e.g. multiplicative, additive) is important in considering risk transfer between different populations.
- To expand research on sex-related differences in the immune and inflammatory responses (e.g., in brain microglia) to low doses and dose-rates, and their role in the induction of cancer and non-cancer effects.

- To develop mechanistic or other mathematical models of radiation-induced disease pathogenesis that can account for individual risk factors.

Research line: Basic mechanisms

Basic research is needed to establish which factors and processes (including genetic, epigenetic and environmental factors/processes, co-morbidities, co-exposures and lifestyle factors) lead to greater individual risk of late effects in terms of cancer or non-cancer diseases. This includes the discovery of genetic, phenotypic and molecular markers of these pathways, and the integration of mechanistic studies in the quantitative evaluation of health risks. A major focus should be the understanding of how these different factors may modify risk, keeping in mind that the radiosensitive phenotype is likely to be multifactorial. Another important question is whether biomarkers of radiation in normal tissue reactions are related to risk of developing late effects following exposure to low and protracted doses of different LETs including internal exposures.

Priority research areas are:

- To develop an understanding of the molecular, cellular, organ and systemic responses determining individual response to radiation-induced health effects including, for example, existing germline and somatic mutations as well as inflammatory processes and immunological states, so that differences between individuals in the response pathways can be predicted, and biomarkers be identified.
- To investigate mechanisms by which age at exposure, attained age, sex, lifestyle and other factors, including co-exposures to other agents and diseases affecting dose from a given exposure, may modulate radiation risk.
- To consider responses from divergent cell types within a tissue or organ, as well as to investigate the impact of anatomical and physiological differences between individuals on radiation dose and dose distributions.
- To start to explore modelling methods, including the use of AI/ML approaches to predict differences in outcome at both individual (qualitative changes affecting health-relevant pathways) and population (quantitative changes in health outcomes) levels.

5.4 Effects of spatial and temporal variation in dose delivery

In the system of radiological protection, risk mainly depends on absorbed dose averaged over a given target mass. The biological outcome of the exposure is determined not only by the dose but also by the time frame of the dose delivery (dose-rate pattern), and by the specific kind of radiation responsible for the energy deposition (radiation quality). In order to account for the effects of temporal variation in dose delivery, a single dose- and dose-rate effectiveness factor (DDREF) is currently applied for low-LET radiation in radiation protection; however, the evidence base for this judgement continues to be debated. Concerning spatial variation, radiation weighting factors are currently applied for radiation protection purposes to account for the difference in the spatial pattern of energy deposition at the subcellular scale, due to different radiation qualities. When it comes to high-LET exposures, particle fluence should be considered in addition to dosimetric information, as fewer cells are expected to be hit but with high deposited energy. At a larger scale, the

effects of intra-organ (but supra-cellular) variation in dose delivery are not considered: the same health risk is assumed for all exposure types if they result in a given amount of absorbed energy, independently of whether the energy is absorbed by a single target cell or homogeneously distributed among all target cells of the same organ. However, the biological effects and so the health consequences are unlikely to be the same.

Inhomogeneity in dose delivery, both at the temporal and spatial level, is a real feature of many environmental, medical and occupational exposure scenarios. Mechanisms responsible for biological effects of different dose-rates or of inhomogeneous dose deposition are not fully characterized: at the cellular level they can be investigated with in vitro studies, but when it comes to how they finally affect health risk (both for cancer and non-cancer diseases) few relevant experimental models or valid datasets exist. In many situations, mixed field exposures are also relevant, but again there are few studies that consider risk in such exposed populations.

The effects of spatial and temporal variation in dose delivery are also gaining importance because of various applications: experiments and trials using extremely high-dose rate (flash) and spatially fractionated therapy beams; the more widespread availability of external beam hadron therapy, where out-of-field doses by scattered neutrons are of concern, the increasing clinical use of radionuclides; and finally the perspective of longer duration space travel (as well as space tourism) in the future. There is also a need to characterize how internal exposure, dose inhomogeneity and radiation quality influence the formation of candidate biomarkers so far identified in response to low-LET external exposure.

Research line: Health risk evaluation

Quantification of health risk at low/moderate dose or dose-rates from internal exposures and from inhomogeneous dose distribution from external exposures is a key challenge for improved radiation risk assessment. As exposures frequently involve all three features noted above (effects of dose rate, radiation quality, and intra-organ dose distribution), relevant cohorts have to be identified or consolidated, where the separate effects of these three variations can be studied. In addition, collection and maintenance of relevant biological sample collections, including tissue samples from cancer cases and somatic tissue from affected individuals may also help to estimate the contribution of the effects of these three exposure characteristics. Sound individual dosimetry is particularly important in cases of internal exposures.

Priority research areas are:

- To determine cancer and non-cancer risks related to acute and chronic internal-emitter exposures in epidemiological studies, incorporating detailed dosimetric assessment and evaluation of dosimetric uncertainties and, where appropriate, microdosimetric considerations.
- To refine risk assessment by integrating analysis of appropriate biological samples and biomarkers of different endpoints and dose to better characterize dose-rate and dose heterogeneity, and their effects. Evaluations of risk must be better targeted to the actual exposure scenarios (for example for radon, high-LET at a protracted exposure), and for that, more experimental data is needed.

- To determine the Relative Biological Effectiveness (RBE) for selected endpoints in epidemiological studies for specific cancer sites through comparison of risk related to low- and high-LET radiation exposure.
- To better determine the risk (as well as possible countermeasures) associated with protracted exposure to the space radiation environment, in view of future interplanetary missions, both for cancer and non-cancer diseases (*e.g.* targeting possible impairments of cognitive and cardiovascular functions).
- To develop and apply more detailed biokinetic and dosimetry models in order to better characterize dose distributions.

Research line: Basic mechanisms

Effects of radiation quality and dose-rate on individual cells and at the cell population level are well documented. Many biological endpoints show a dose-rate dependence (notably DNA damage response) and data supporting an inverse dose-rate effect also exist. This raises the question of the effects of protracted exposures, particularly at low dose and low dose-rate. It is recommended to consider fluence (in addition to dosimetric information) when dealing with exposures to charged particles (particularly for high-LET). Concerning spatial variation at the sub-cellular level, the biological outcome is clearly modulated by radiation quality indicators such as LET. Using *e.g.* microbeam irradiations, mechanisms determining the response to a highly inhomogeneous energy deposition can be addressed under controlled conditions.

To provide further insights in the effects of intra-organ dose distribution, experiments with organotypic tissue models and animal models are required. The effects of locally high doses, when small parts of the tissue/organ are irradiated with high doses while the average dose remains low, have to be quantified and compared to homogeneous exposures. Whether and how effects of the locally high dose propagate in the less exposed tissue also deserves investigation. Organotypic tissue models and animal models also allow the study of the changes in tissue architecture in order to analyse the effects of intra-organ dose distribution.

Priority research areas are:

- To conduct experimental studies *in vitro* and *in vivo* to test exposure scenarios where dose/fluence modulation plays a role, *e.g.* localized versus uniform exposures, acute versus protracted exposures, to inform specific biomarker development and risk quantification.
- To further develop suitable tissue and *in vivo* models for the quantification of the impact of dose inhomogeneity and radiation quality.
- When addressing the effects of internal contamination, specifically consider the role of chemical speciation in determining spatial distribution (at all scales) and biokinetics of radionuclides, as well as in contributing to the risk (co-exposure).
- For all adopted experimental models, to develop in parallel computational modelling approaches able to tackle and quantify inhomogeneity at all scales: nano- (radiation track structure) and microdosimetric, dosimetric and biokinetic models at different levels of biological organisation.

- To study mechanisms elicited by inhomogeneous dose deposition, integrating “dynamic” dose assessment and the dynamic response of the biological target (e.g. at the genome level, at the cell population level via signalling or cell motility, at the organismal level also via the immune response) and to finally identify relevant pathways (both for cancer and non-cancer diseases) in a systems biology approach, in order to characterize the response of the complex system as a whole.
- To develop innovative ways in experimental studies including organotypic tissue models to determine the Relative Biological Effectiveness (RBE) at low doses to determine/compare the effects of low- versus high-LET exposure. To characterize how internal exposure, dose inhomogeneity and radiation quality will affect the nature of candidate biomarkers so far identified in response to low-LET external exposure.
- To develop experimental and modelling strategies to characterize the effects of exposures to mixed fields.
- Build on knowledge acquired from basic mechanisms to identify relevant pathways for the quantification of the risk for cancer and non-cancer diseases, also using an AOP approach, determining those operating in cases of inhomogeneous exposures.
- Elucidate mechanisms of out-of-field effects, including systemic inflammatory responses and intercellular communication pathways.

6. Education and Training

6.1 The role of education and training in low-dose radiation research

The [HLEG Report of 2009](#) identified a problem with the maintenance in Europe of the range of expertise essential to an effective programme of research into the risks to humans from low-dose radiation. The report advises that specific programmes aiming at knowledge management across generations have to be designed in order to achieve sustainable continuity and development.

A large proportion of the groundwork of research is carried out as student projects and thesis work. For this reason, the research effort relies on a continuing relationship with universities, and on a healthy stream of high-level students. It is essential that this symbiosis is recognised and taken into account in research funding structures.

A further intrinsic role of E&T within any specialized research area is in dissemination of new technologies, skills, and knowledge. To obtain maximum impact and benefit from research there should be an actively managed programme of workshops, seminars, summer schools, student exchange programs etc. which is integrated into the design and funding structure of all research. The programme should be aimed both at the sharing of knowledge within the European low-dose research community and also in the wider radiation protection field including radioecology, emergency response, and the medical use of radiation.

6.2 Priorities for strategic support of E&T

Following the comments in the previous section, support for E&T has two priority areas: support for students and young scientists, and promotion of E&T for dissemination.

Support for students and young scientists

- Students need to be able to find places at universities and research groups for project/dissertation work. This requires that the places be available, but also that there are sufficient incentives to attract students. Universities are autonomous and develop new courses in response to a perceived need, taking account of staff expertise and specialization. Financial support from outside is not needed to achieve this end, although there is a role for influencing the perceived need. On the other hand, increasing the access of students Europe-wide to university courses through industry-funded scholarships could significantly help to attract students. Setting up such a post-graduate scholarship scheme for attendance at approved universities should be seen as a priority. Coordination with existing schemes (e.g. EU Marie Curie, IAEA Marie Skłodowska-Curie Fellowship Programme) could be useful in this respect.
- In order to complement support at the post-graduate level and to help provide a career path for the most promising graduates, a scheme for provision of one or more post-doctoral fellowships should also be offered, to be taken up at approved research institutions.
- The MELODI Mobility Programme is established and offers travel award support to early career scientists, PhD and MSc students, as the groups identified with greatest need for this form of support. The intention of the MELODI Mobility Programme is to financially support participation in conferences, courses, visits, internships or enable a student exchange to carry out scientific research, all in order to increase the applicant's involvement and knowledge/skills in European research in radiation protection.

Promotion of E&T for dissemination

- It should be explicit in the wording for RTD calls that proposals will be judged favourably if a plan is included that explains how E&T will be integrated into the overall research programme, providing workshops or training courses dedicated to the presentation of new science/technology which is being used or developed in the project.
- Parallel to the E&T supported by the RTD calls, it is seen as essential that a separately funded body (or part of a body with a ring-fenced budget) is responsible for the organization and sponsorship of targeted initiatives in order to promote the specialized skills and knowledge needed to maintain the full competence of the low-dose research community. These will be made readily available to postgraduate students and scientists. The benefit to the former will be the provision of supplements to their university courses and the opportunity to gain experience of the different areas of science on offer to them in their future careers. For the latter, this will be a very

effective way of providing continuing professional education, and of sharing knowledge with other research and educational institutions.

Coordination and collaboration of E&T providers

In order to get maximum benefit from E&T in the low-dose research area (both that which is already provided and the new initiatives proposed here) there should be an overall coordination of resources within the European community. Recommended priority actions are as follows:

- Continuation and extension of the MELODI Education and Training Forum in order to bring together all platforms and other interested parties regularly to discuss needs and broaden the awareness of what is happening in EU member states. This should be seen as both a problem-solving and an advertising forum. There should be active participation by all other platforms involved in radiation protection (ALLIANCE, NERIS, EURADOS, EUTERP, EURAMED etc.) in order to share mutual experience and resources.
- There should be active cooperation among groups promoting and supporting E&T in the radiation protection and research area (EURAYS, ENEN, etc.) and possibly the use of mailing lists or social media to advertise programmes, courses, scholarships, fellowships, etc.

7. Infrastructures

One of the roles of MELODI is to promote and facilitate access to the state-of-the-art research infrastructures to support the research efforts in the radiation protection field which are prioritized in this SRA. In order to identify, characterise and quantify health risks accurately, the quality of raw data and final results produced from research projects is essential. So, the harmonization of practices amongst multiple facilities is becoming an increasingly important indicator of reliability and finally, of the sustainability of the virtual network of infrastructures as well as a guarantee of the dynamism and high quality of the research area.

Infrastructures include exposure facilities, observatories as well as databases (including cohorts) and biobanks and many analytical platforms (e.g. for omics) and specific tools especially developed for the domain.

Within DoReMi, a first extensive list of relevant infrastructures was generated for low dose research, in particular irradiation facilities for internal and external exposure. Within projects OPERRA and more recently CONCERT, infrastructures were highlighted via AIR² bulletin and AIR²D² database. A web-handbook has been produced, describing exposure facilities, cohorts, databases, biobanks and analytical platforms.

STORE, an internet-based platform for storage and sharing data from previous studies, has been developed and continues to grow. Going forward, it will be necessary to promote activities to maintain and develop this database and continuously expand it as STORE also includes systematically all new data and results arising from Euratom/Horizon Europe-supported projects. The use of this repository for data linked to all publications arising

from funded projects in radioprotection research should be required, where appropriate and possible (ethics requirements and informed consent in epidemiological studies), while adhering to FAIR principles and the applicable Horizon Europe rules. Particular attention has to be dedicated to aspects related to sensitive data management. The implementation in 2018 of the GDPR (679/16) requires efforts and a dedicated budget to streamline compliance with GDPR rules.

In order to assess the most appropriate infrastructures that meet the needs of the various priorities described in this SRA and so the future specific linked projects, it is necessary to develop and apply quality criteria for selection through the future calls. Financial support for those infrastructures can be included in future calls in which they are clearly identified answering to specific services under guarantees such as protocols, criteria about harmonization and quality of results, data storage and sharing.

The use of bio-banked materials, where applicable, should be encouraged by including their use in future calls either indirectly for all relevant proposals or by specific topics dedicated to their use. In addition, funding should be included to support the biobanking of samples arising from Euratom/Horizon Europe funded projects where appropriate.

Next steps will rely on further harmonisation of quality standards, practices and protocols, and co-operation between the European radiation protection research platforms in relation to the provision and use of infrastructure. Huge efforts will be dedicated to sample/data acquisition and sample/data storage with the aim of reuse archived materials. There is a need for a transnational agreement on a strategic work plan for maintenance, updating, mutual use of suitable infrastructures.

Simultaneously, education and training actions should be developed to promote the use of existing powerful European research infrastructures rather than local and inadequate ones. The advantages of using these relevant infrastructures under common rules for transnational access should be clear and should provide an incentive for their use.

Priority areas are:

- Develop easy access and improve the common organisation of the existing network of infrastructures, using feedback from approaches applied for infrastructures networks arising from past initiatives within Europe.
- Improve the awareness of existing relevant infrastructures through E&T courses and promote their use by implementing on-site practical courses.
- Develop protocols and guidance documents, approaching a common compliance with GDPR rules, with data management (storage and sharing) applying FAIR principles, favour open access within STORE and promote the re-use of archived materials and retrospective approaches using existing epidemiological data.
- Develop inter-comparisons and harmonization activities to guarantee the quality of data and results arising from funded projects.

8. Abbreviations, Websites

ALLIANCE (European Radioecology Alliance) <http://www.er-alliance.org/>

DoReMi Network of Excellence (Low Dose Research towards Multidisciplinary Integration), <https://melodi-online.eu/doremi/>

CONCERT <https://concert-h2020.eu>

EURADOS (The European Radiation Dosimetry Group) www.eurados.org/

EURAMED (European Alliance for Medical Radiation Protection Research) <http://www.eibir.org/scientific-activities/joint-initiatives/european-alliance-for-medicalradiation-protection-research-euramed/>

HLEG (High Level expert group) <https://op.europa.eu/en/publication-detail/-/publication/47c88945-aec3-4730-ad75-e1f9173e5c09>

MEDIRAD (Implications of Medical Low Dose Radiation Exposure) <http://www.medirad-project.eu/>

MEENAS (MELODI, EURADOS, EURAMED, NERIS, ALLIANCE, SHARE) an organisation coordinating the work of all European radiation protection research platforms.

MELODI (Multidisciplinary European Low Dose Initiative) <http://www.melodi-online.eu/>

NASEM (National Academies of Sciences, Engineering and Medicine, USA) report on low dose research: [Leveraging advances in modern science to revitalize low dose radiation research in the United States](#)

NERIS (European Platform on preparedness for nuclear and radiological emergency response and recovery) <http://www.eu-neris.net/>

OPERRA (Open project for European Radiation Research Area) <http://www.melodi-online.eu/operra.html>

SHARE (platform on social sciences and humanities) <http://sites.exeter.ac.uk/nuclearsocieties/shine/>

STORE (platform for the archiving and sharing of the primary data outputs from research on low dose radiation) <https://www.storedb.org>

9. Annex: Publications from MELODI Workshops

Individual radiosensitivity workshop

Salomaa, S. and Jung, T. (2020) 'Roadmap for research on individual radiosensitivity and radiosusceptibility - the MELODI view on research needs', *International Journal of Radiation Biology*, 96(3), pp. 277-279. doi:10.1080/09553002.2019.1704107.

Gomolka, M., Blyth, B., Bourguignon, M., Badie, C., Schmitz, A., Talbot, C., Hoeschen, C. and Salomaa, S. (2020) 'Potential screening assays for individual radiation sensitivity and susceptibility and their current validation state', *International Journal of Radiation Biology*, 96(3), pp. 280-296. doi:10.1080/09553002.2019.1642544.

Averbeck, D. et al. (2020) 'Establishing mechanisms affecting the individual response to ionizing radiation', *International Journal of Radiation Biology*, 96(3), pp. 297-323. doi:10.1080/09553002.2019.1704908.

Seibold, P. et al. (2020) 'Clinical and epidemiological observations on individual radiation sensitivity and susceptibility', *International Journal of Radiation Biology*, 96(3), pp. 324-339. doi:10.1080/09553002.2019.1665209.

Kalman, C. and Oughton, D. (2020) 'Ethical considerations related to radiosensitivity and radiosusceptibility', *International Journal of Radiation Biology*, 96(3), pp. 340-343. doi:10.1080/09553002.2019.1665210.

Non-cancer effects workshop

Kreuzer, M. and Bouffler, S. (2021) 'Guest editorial: Non-cancer effects of ionizing radiation - clinical implications, epidemiological and mechanistic evidence and research gaps', *Environment International*, 149, 106286. doi:10.1016/j.envint.2020.106286.

Ainsbury, E.A. et al. (2021) 'Radiation-induced lens opacities: Epidemiological, clinical and experimental evidence, methodological issues, research gaps and strategy', *Environment International*, 146, 106213. doi:10.1016/j.envint.2020.106213.

Lumniczky, K. et al. (2021) 'Low dose ionizing radiation effects on the immune system', *Environment International*, 149, 106212. doi:10.1016/j.envint.2020.106212.

Pasqual, E. et al. (2021) 'Cognitive effects of low dose of ionizing radiation - Lessons learned and research gaps from epidemiological and biological studies', *Environment International*, 147, 106295. doi:10.1016/j.envint.2020.106295.

Tapio, S. et al. (2021) 'Ionizing radiation-induced circulatory and metabolic diseases', *Environment International*, 146, 106235. doi:10.1016/j.envint.2020.106235.

Spatial and temporal variation in dose delivery workshop

Madas, B.G. and Wojcik, A. (2022) 'The 2020 MELODI workshop on the effects of spatial and temporal variation in dose delivery', *Radiation and Environmental Biophysics*, 61, pp. 479-483. doi:10.1007/s00411-022-01002-3.

Pazzaglia, S. et al. (2022) 'Out-of-field effects: lessons learned from partial body exposure', *Radiation and Environmental Biophysics*, 61, pp. 485-504. doi:10.1007/s00411-022-00988-0.

- Lowe, D. et al. (2022) 'Radiation dose rate effects: what is new and what is needed?', *Radiation and Environmental Biophysics*, 61, pp. 507-543. doi:10.1007/s00411-022-00996-0.
- Baiocco, G. et al. (2022) 'A matter of space: how the spatial heterogeneity in energy deposition determines the biological outcome of radiation exposure', *Radiation and Environmental Biophysics*, 61, pp. 545-559. doi:10.1007/s00411-022-00989-z.
- Madas, B.G. et al. (2022) 'Effects of spatial variation in dose delivery: what can we learn from radon-related lung cancer studies?', *Radiation and Environmental Biophysics*, 61, pp. 561-577. doi:10.1007/s00411-022-00998-y.
- Li, W.B. et al. (2022) 'Heterogeneity of dose distribution in normal tissues in case of radiopharmaceutical therapy with alpha-emitting radionuclides', *Radiation and Environmental Biophysics*, 61, pp. 579-596. doi:10.1007/s00411-022-01000-5.
- Saldarriaga Vargas, C. et al. (2024) 'Heterogeneity of absorbed dose distribution in kidney tissues and dose-response modelling of nephrotoxicity in radiopharmaceutical therapy with beta-particle emitters: A review', *Zeitschrift für Medizinische Physik*, 34(4), pp. 491–509. doi:10.1016/j.zemedi.2023.02.006.

Adverse Outcome Pathways workshop

- Chauhan, V. et al. (2022a) 'Introduction to the special issue on adverse outcome pathways in radiation protection', *International Journal of Radiation Biology*, 98(12), pp. 1691-1693. doi:10.1080/09553002.2022.2123183.
- Chauhan, V. et al. (2022b) 'Adverse outcome pathway: a path toward better data consolidation and global co-ordination of radiation research', *International Journal of Radiation Biology*, 98(12), pp. 1694-1703. doi:10.1080/09553002.2021.2020363.
- Chauhan, V. et al. (2022c) 'A high-level overview of the OECD AOP Development Programme', *International Journal of Radiation Biology*, 98(12), pp. 1704-1713. doi:10.1080/09553002.2022.2110311.
- Chauhan, V. et al. (2022d) 'Establishing a communication and engagement strategy to facilitate the adoption of the adverse outcome pathways in radiation research and regulation', *International Journal of Radiation Biology*, 98(12), pp. 1714-1721. doi:10.1080/09553002.2022.2086716.
- Azimzadeh, O. et al. (2022) 'Application of radiation omics in the development of adverse outcome pathway networks: an example of radiation-induced cardiovascular disease', *International Journal of Radiation Biology*, 98(12), pp. 1722-1751. doi:10.1080/09553002.2022.2110325.
- Jaylet, T. et al. (2022) 'Development of an adverse outcome pathway for radiation-induced microcephaly via expert consultation and machine learning', *International Journal of Radiation Biology*, 98(12), pp. 1752-1762. doi:10.1080/09553002.2022.2110312.
- Burt, J.J. et al. (2022) 'Radiation adverse outcome pathways (AOPs) are on the horizon: advancing radiation protection through an international Horizon-Style exercise',

International Journal of Radiation Biology, 98(12), pp. 1763-1776.
doi:10.1080/09553002.2022.2121439.

Kozbenko, T. et al. (2022) 'Deploying elements of scoping review methods for adverse outcome pathway development: a space travel case example', *International Journal of Radiation Biology*, 98(12), pp. 1777-1788. doi:10.1080/09553002.2022.2110306.

Yu, J. et al. (2022) 'Adverse outcome pathways and linkages to transcriptomic effects relevant to ionizing radiation injury', *International Journal of Radiation Biology*, 98(12), pp. 1789-1801. doi:10.1080/09553002.2022.2110313.

Klokov, D. et al. (2022) 'International expert group collaboration for developing an adverse outcome pathway for radiation induced leukemia', *International Journal of Radiation Biology*, 98(12), pp. 1802-1815. doi:10.1080/09553002.2022.2117873.

Tollefsen, K.E. et al. (2022) 'Adverse outcome pathways (AOPs) for radiation-induced reproductive effects in environmental species: state of science and identification of a consensus AOP network', *International Journal of Radiation Biology*, 98(12), pp. 1816-1831. doi:10.1080/09553002.2022.2110317.

Stainforth, R. et al. (2022) 'Benchmark dose modeling of transcriptional data: a systematic approach to identify best practices for study designs used in radiation research', *International Journal of Radiation Biology*, 98(12), pp. 1832-1844.
doi:10.1080/09553002.2022.2110300.

Adam, N. et al. (2022) 'Evaluating the influences of confounding variables on benchmark dose using a case study in the field of ionizing radiation', *International Journal of Radiation Biology*, 98(12), pp. 1845-1855. doi:10.1080/09553002.2022.2110303.

Chauhan, V. et al. (2021) 'Expert consultation is vital for adverse outcome pathway development: a case example of cardiovascular effects of ionizing radiation', *International Journal of Radiation Biology*, 97, pp. 1516-1525. doi:10.1080/09553002.2021.1969466.

Offspring and next generations workshop

Streffer, C. and Hande, M.P. (2024) 'Effects of ionizing radiation exposure in offspring and next generations', *International Journal of Radiation Biology*, 100(9), pp. 1237-1239. doi:10.1080/09553002.2024.2384834.

Amrenova, A. et al. (2024a) 'Consideration of hereditary effects in the radiological protection system: evolution and current status', *International Journal of Radiation Biology*, 100(9), pp. 1240-1252. doi:10.1080/09553002.2023.2295289.

Amrenova, A. et al. (2024b) 'Intergenerational effects of ionizing radiation: review of recent studies from human data (2018-2021)', *International Journal of Radiation Biology*, 100(9), pp. 1253-1263. doi:10.1080/09553002.2024.2309917.

Benotmane, M.A. and Trott, K.R. (2024) 'Epidemiological and experimental evidence for radiation-induced health effects in the progeny after exposure in utero', *International Journal of Radiation Biology*, 100(9), pp. 1264-1275.
doi:10.1080/09553002.2023.2283088.

Degenhardt, Ä., Dumit, S. and Giussani, A. (2024a) 'Effects of ionising radiation exposure in offspring and next generations: dosimetric aspects and uncertainties', *International Journal of Radiation Biology*, 100(9), pp. 1276-1282. doi:10.1080/09553002.2023.2280017.

Grisson, S. et al. (2024) 'In utero exposure to ionizing radiation and metabolic regulation: perspectives for future multi- and trans-generation effects studies', *International Journal of Radiation Biology*, 100(9), pp. 1283-1296. doi:10.1080/09553002.2023.2295293.

Nakamura, N., Yoshida, N. and Suwa, T. (2024) 'Three major reasons why transgenerational effects of radiation are difficult to detect in humans', *International Journal of Radiation Biology*, 100(9), pp. 1297-1311. doi:10.1080/09553002.2023.2187478.

Sreetharan, S. et al. (2024) 'Ionizing radiation exposure effects across multiple generations: evidence and lessons from non-human biota', *International Journal of Radiation Biology*, 100(9), pp. 1312-1329. doi:10.1080/09553002.2023.2281512.

Stephens, J. et al. (2024) 'A systematic review of human evidence for the intergenerational effects of exposure to ionizing radiation', *International Journal of Radiation Biology*, 100(9), pp. 1330-1363. doi:10.1080/09553002.2024.2306328.

Liutsko, L. et al. (2024) 'Association between radiation dose, thyroid hormone, and IQ levels in children exposed to radiation in utero after the Chernobyl accident', *International Journal of Radiation Biology*, 100(9), pp. 1364-1370. doi:10.1080/09553002.2024.2345088.

Zölzer, F. et al. (2024) 'Ethical and societal aspects of radiological protection for offspring and next generations', *International Journal of Radiation Biology*, 100(9), pp. 1371-1381. doi:10.1080/09553002.2023.2281523.

Degenhardt, Ä. et al. (2024b) 'The ICRP, MELODI, and ALLIANCE workshop on effects of ionizing radiation exposure in offspring and next generations: a summary of discussions', *International Journal of Radiation Biology*, 100(9), pp. 1382-1392. doi:10.1080/09553002.2024.2306335.

Radiation-induced circulatory diseases workshop

Hamada, N. (2023) 'MELODI Workshop 2023 "Updates on Radiation-Induced Circulatory Diseases"', *Japanese Journal of Health Physics*, 58(3), pp. 182-193. doi:10.5453/jhps.58.182.

Central nervous system workshop

Hamada, N., Bernier, M.-O., Lumniczky, K. and Laurier, D. (2026) 'Health effects of ionizing radiation exposure in the brain: MELODI perspectives on potential implications of emerging evidence for radiation protection', *Mutation Research - Reviews in Mutation Research*, 797, 108582. doi:10.1016/j.mrrev.2025.108582.

Nakamizo, T. (2025) 'MELODI Topical Workshop 2025 "Radiation Effects in the Central Nervous System": A Conference Report', *Japanese Journal of Health Physics*, 60(3), pp. 257-264. doi:10.5453/jhps.60.257.